


**PONV-03 & PONV-03b: Postoperative Nausea or Vomiting Outcome**

**PONV-03:** (Inverse Measure) Percentage of patients, regardless of age, who undergo a procedure and have a documented nausea/emesis occurrence OR receive a rescue antiemetic in the immediate postoperative period.

**PONV-03b:** (Inverse Measure) Percentage of patients, regardless of age, who undergo a procedure and have a documented nausea/emesis occurrence with or without receiving an antiemetic in the immediate postoperative period.

**Measure Time Period:**  
**PACU Start** to 6 hours after **Anesthesia End**  
 PACU Start determined by **PACU Start Phenotype**  
 Anesthesia End determine by **Anesthesia End Phenotype**



Anti-emetic agents by class	
<b>5-Hydroxytryptamine (5-HT3)</b> 10335 Ondansetron 10164 Dolasetron 10208 Granisetron 10711 Palonosetron	<b>Butyrophenones</b> 10169 Droperidol 10210 Haloperidol
<b>Benzamides</b> 10820 Amisulpride	<b>Neurokinin-1 Receptor Agonists</b> 10035 Aprepitant 10719 Fosaprepitant
<b>Antihistamines</b> 10257 Dimenhydrinate 10160 Diphenhydramine	<b>Phenothiazines</b> 10374 Promethazine 10373 Prochlorperazine
<b>Prokinetic</b> 10297 Metoclopramide	