


**PONV-05: PONV Prophylaxis for Adult Patients**

**PONV-05:** Percentage of patients who had a procedure requiring general anesthesia OR cesarean delivery and received appropriate prophylaxis for postoperative nausea and vomiting.



- Excluded Procedures:**
- **Bronchoscopy**
  - **Diagnostic Imaging**
  - **ECT**
  - **Endoscopy**
  - **Intubation Only**
  - Labor Epidurals determined by **Obstetric Anesthesia Type** (value codes 3, 5 & 6 including obstetric non-operative procedures)
  - **TEE/Cardioversion**

- PONV Risk Factors**
- Age < 50 years
  - Sex: Female
  - History of PONV or motion sickness (equates to one risk factor even if history of both)
  - Non-smoker
  - Opioid use for postoperative given intraoperatively or postoperatively
  - Duration of inhalational anesthesia greater than 1 hour
  - Procedure Type:
    - Cholecystectomy
    - Laparoscopy
    - Gynecologic

Antiemetic agents by class	
<b>5-Hydroxytryptamine (5-HT3) *</b> 10335 Ondansetron 10208 Granisetron 10711 Palonosetron	<b>Antidopaminergics*</b> 10169 Droperidol 10210 Haloperidol 10297 Metoclopramide 10820 Amisulpride
<b>Anticholinergics</b> 10400 Scopolamine Patch 10399 Scopolamine 11040 Butylscopolamine	<b>Neurokinin-1 Receptor Agonists</b> 10035 Aprepitant 10719 Fosaprepitant
<b>Antihistamines</b> 10257 Dimenhydrinate 10160 Diphenhydramine 10635 Meclizine	<b>Phenothiazines</b> 10374 Promethazine 10373 Prochlorperazine
<b>Glucocorticoids*</b> 10147 Dexamethasone 10296 Methylprednisolone	<b>Other</b> Propofol (Infusion only)
*Acceptable for cesarean delivery	

**Only antiemetics given via the following routes will be considered:**

MPOG Concept ID	Route
2001	IV
2003	Intramuscular
2008	Oral
2011	Sublingual
2012	Rectal
2022	Intraosseous
2023	Enteric Tube
2997	Other – Not Listed
2998	Other – Not Specified
2999	Other

For Scopolamine only: Topical (2013) or Transdermal (2027)